



**EFFECTS OF LOW FLOW  
ON FOUNTAIN DARTER  
FECUNDITY STUDY**


**Literature Review and Proposed Methodology**

**PREPARED FOR:**

**HCP SCIENCE COMMITTEE**

**PREPARED BY:**

**BIO-WEST PROJECT TEAM - February 18, 2014**



## INTRODUCTION

Reproductive success of slackwater and benthic fishes is reduced under low flow conditions, attributed to greater variability in physical habitats and to increases in organic substrates (Schlosser 1982, Falke et al. 2010). As flows decrease, aquatic vegetation (a physical habitat for fishes) proliferates but not homogeneously among plant taxa (Bunn and Arthington 2002, Riis and Hawes 2002). Therefore, changes in the plant community (i.e., low-growing/tall-growing, sparse/dense macrophytes) and accumulation of organic sediments under a declining hydrograph can limit spawning and nursery habitats of stream fishes, especially those that attached eggs to plants or substrates. Habitat association of the fountain darter is primarily benthic slackwater and low velocity habitats. Fountain darters inhabit no to dense vegetative cover in shallow and deep water habitats but associate with low growing vegetation (Schenck and Whiteside 1977, Alexander and Phillips 2012), which likely optimizes consumption of drifting and benthic prey and reproductive habitat while avoiding predators (Bergin 1996, Brandt et al. 1993, Phillips et al. 2011, Dammeyer et al. 2013). Given that low flow conditions will alter the physical habitats of the fountain darter, we predict that changes in physical habitats, especially low-growing and dense vegetation, will reduce the reproductive readiness and success of the fountain darter.

To test this prediction, we propose to establish a baseline in fountain darter reproductive readiness among a gradient of flow regimes and among vegetation type. Objectives of this study are to quantify elements of fountain darter reproduction (gonadal recrudescence, ovarian development, fecundity, and oocyte maturation) among available flow gradients ranging from 10 to 120 cfs in the wild and among physical habitat types and substrates (open substrates, low-growing and tall-growing aquatic vegetation).

### Benefits to HCP Ecological Model

We are directly assessing the influence of flow and influence of vegetation on fountain darter reproduction. Flow and amount of vegetation are two key components of fountain darter habitat in the HCP Ecological Model, and information generated from this work will provide more accurate estimations on model parameters.

## LITERATURE REVIEW


**Baseline reproductive rates of fountain darters:** Fountain darters are sexually dimorphic with males having distinct coloration in dorsal fins and short, pointed genital papillae and females having less intense pigmentation in dorsal fins and long, forked genital papillae (Schenck and Whiteside 1977). Sex ratios are slightly skewed toward males (1.39:1:00). Minimum length of reproduction is 24 mm in total length (Schenck and Whiteside 1977) at age 3.5 months (Linam et al. 1993) to 6 months (Brandt et al. 1993). Numbers of ova (ovulated oocytes within the ovary) are related to female length in darters with larger females producing more ova, though size of ova is independent of female length (Schenck and Whiteside 1977, Marsh 1986). Ova occur in female fountain darters year round, suggesting a protracted spawning season (12-months) but with reproductive peaks in late winter and late summer (Schenck and Whiteside 1977). Fountain darters are batch spawners, producing a mean of 9 to 14.5 eggs per day during a 33 d period in a hatchery setting (Bonner et al. 1998) with 5 to 27 days, on average, between batches (Brandt et al. 1993). Eggs are released at water temperatures ranging from 3 to 30°C (Brandt et al. 1993) with optimum egg production ranging between 14 and <26° (Bonner et al. 1998, McDonald et al. 2007).

**Habitats used by fountain darters for egg deposition:** Fountain darters are facultative phytophilic spawners (Simon 1999) depositing adhesive eggs on macrophytes (Strawn 1956, Phillips et al. 2011) but also on hard substrates lacking vegetation (Brandt et al. 1993). To date, fountain darter eggs occur on *Rhizoclonium*, *Ludwigia*, *Sagittaria*, and *Zizania* (Phillips et al., 2011), but this is likely an incomplete list. Fountain darters associate with a wide variety of vegetation, including *Riccia*, *Rhizoclonium*, *Hydrilla*, *Ludwigia*, *Potamogeton*, *Sagittaria*, *Vallisneria*, *Hygrophilia*, and *Combomba* (Schenck and Whiteside 1976, Linam et al. 1993, Phillips et al. 2011, Alexander and Phillips 2012, Araujo 2012, Dammeyer et al. 2013) and areas without vegetation (Crowe and Sharp 1997, Araujo 2012, Behen 2013). Fountain darters, in general, associate with slackwater and low velocity habitats, ranging in depths from < 0.5 m to 5 m with silt to cobble substrates (Behen 2013). Sister species within Subgenus *Microperca* (*E. microperca* and *E. proeliare*; Near et al. 2011) also are associated with slackwater to run habitats consisting of detrital terrestrial leaves, woody debris, and dense vegetation (Burr and Page 1978; Paine et al. 1981; Johnson and Hatch 1991).

**Potential threats to fountain darter reproduction related to low flow conditions:** Reduction in base flow reduces the amount of available habitat for spring-associated fishes (Hubbs 1995), and likely fragments habitats and impede movement (Dammeyer et al. 2013), decreases fountain darter reproductive success (Brandt et al. 1993, Bonner et al. 1998), and increases intraspecific competition (Araujo 2012) and gill-parasite mortality (McDonald et al. 2006, Tolley-Jordan and Owen 2008). Modeling suggests that reducing the 19-year mean base flow conditions (184 cfs) to 58 cfs (32% of current base flow) would noticeably reduce fountain darter populations in the San Marcos River (Mora et al. 2013). Fountain darters were considered extirpated from the Comal River in 1973, attributed to cessation of spring flows and a rotenone treatment to remove non-native fishes in the 1950s, a catastrophic flood in 1972, or a combination of the three events (Schenck and Whiteside 1976).

## PROPOSED METHODS

As stated in the introduction, objectives of this study are to quantify elements of fountain darter reproduction among available flow gradients ranging from low flow (10 to 30 cfs; Upper Spring Run, Landa Lake), moderate flow (50 to 60 cfs; Old Channel-Comal River, and 80 to 100 cfs; new channel-Comal River), and high flow (100 to 120 cfs; San Marcos River) conditions and among physical habitat types and substrates (open substrates, low-growing and tall-growing aquatic vegetation). Study results should yield the following information: reproductive readiness of fountain darters among base flows ranging from 10 to 120 cfs and among substrates ranging in vertical depth from short (gravel and cobbles, no vegetation) to tall-growing vegetation. An example of potential results is presented in the following figure.



Sexually mature (>24 mm) female fountain darters will be harvested monthly between February and September 2014. On each collection trip and when available, up to 5 female darters will be taken from 3 to 5 areas of bare substrates (sand, gravel, cobble), short growing vegetation, and tall-growing vegetation within the San Marcos River and Comal River. Up to 60 female fountain darters will be taken on each collection trip. Fish will be anesthetized with MS-222 and preserved in 10% buffered formalin. In the laboratory, lengths and weights of each fish will be taken. Gonadosomatic index, ovarian stage, fecundity, and oocyte maturation will be determined for each fish following methodologies described specifically for darters by Heins and Baker (1989), Heins et al. (1992), and Heins (1995).

Monthly collections are necessary to quantify seasonality in fountain darter reproduction. In part, this work will support or revise spawning periodicity published by Schenck and Whiteside (1977). A more

refined methodology will be used to assess oocyte maturity (Heins and Baker 1989) than used by Schenck and Whiteside (1977). Also, monthly gonadosomatic indices might be used as a covariate in the experimental design of this study.

### **Experimental Design and Data Analysis**

Response variables: Gonadosomatic index, number of ova, percent of mature ovaries

Experimental unit: individual female fountain darter

Treatment 1: Flow regime (four levels)

Treatment 2: Substrate type (three levels)

Treatment 3 (likely co-variate): Month (8 levels)

$$8 \times 4 \times 3 = 96$$

A minimum of five replications per treatment were selected to control variability relating to batch spawning fishes ( $5 \times 96 = 480$  female fountain darters). A multi-factored MANOVA will be used to assess differences ( $\alpha=0.05$ ) among treatment effects and response variables.

### **LITERATURE CITED**

- Alexander, M. L. and C. T. Phillips. 2012. Habitats used by the endangered Fountain Darter (*Etheosoma fonticola*) in the San Marcos River, Hays County, Texas. *The Southwestern Naturalist* 57:449-452.
- Araujo, D. 2012. Effect of drought and subsequent recovery on endangered fountain darter habitat in Comal springs, Texas. MS Thesis. Texas State University.
- Behen, K. P. K. 2013. Influence of connectivity and habitat heterogeneity on fishes in the upper San Marcos River, Texas. MS Thesis. Texas State University.
- Bergin, J.S. 1996. Diet of the fountain darter, *Etheostoma fonticola*, in the Comal River, Texas MS Thesis. Texas State University.
- Bonner, T. H., Brandt, T. M., Fries, J. N., and Whiteside, B. G. 1998. Effects of temperature on egg production and early life stages of the fountain darter. *Transactions of the American Fisheries Society* 127: 971-978.
- Brandt T.M., K.G. Graves, C.S. Berkhouse, T.P. Simon, and B.G. Whiteside. 1993. Laboratory spawning and rearing of the endangered fountain darter. *Progressive Fish-Culturist* 55:149-156.
- Bunn, S. E. and A. H. Arthington. 2002. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management* 30:492-507.
- Burr, B. M. and L. M. Page. 1978. The life history of the Cypress Darter, *Etheostoma proeliare*, in Max Creek, Illinois. *Illinois Natural History Survey, Biological Notes* 106:1-15.
- Crowe, J. C. and J. M. Sharp, Jr. 1997. Hydrogeologic delineation of habitats for endangered species: the Comal Springs/River System. *Environmental Geology* 30:17-28.

- Dammeyer, N. T., C. T. Phillips, and T. H. Bonner. 2013. Site fidelity and movement of the smallest etheostomine darter with implications for endangered species management. *Transactions of the American Fisheries Society* 142:1049-1057.
- Falke, J. A., K. R. Bestgen, and K. D. Fausch. 2010. Streamflow reductions and habitat drying affect growth, survival, and recruitment of Brassy Minnow across a Great Plains Riverscape. *Transactions of the American Fisheries Society* 139:566-1583.
- Heins, D. C. 1995. Techniques for the determination of clutch parameters in darters and minnows: a proposal for standard methods. Technical Paper 1. Online only: <http://tulane.edu/sse/eebio/research/resources/technical-paper1.cfm>
- Heins, D. C., and J. A. Baker. 1989. Growth, population structure, and reproduction of the percid fish *Percina vigil*. *Copeia* 1989: 727-736.
- Heins, D. C., J. A. Baker, and W. P. Dunlap. 1992. Yolk loading in oocytes of darters and its consequences for life-history study. *Copeia* 1992: 404-412.
- Johnson, J. D. and J. T. Hatch. 1991. Life history of the Least Darter *Etheostoma microperca* at the northwest limits of its range. *American Midland Naturalist* 125:87-103.
- Linam, G.W, K. B. Mayes, K. S. Saunders. 1993. Habitat utilization and population size estimate of fountain darters, *Etheostoma fonticola*, in the Comal River, Texas. *Texas Journal of Science* 45: 341-348
- Marsh, E. 1986. Effects of egg size on offspring fitness and maternal fecundity in the orangethroat darter, *Etheostoma spectabile* (Pisces: Percidae). *Copeia* 1: 18-30.
- McDonald, D. L., T. H. Bonner, T. M. Brandt, and G. H. Trevino. 2006. Size susceptibility to trematode-induced mortality in the endangered fountain darter (*Etheostoma fonticola*). *Journal of Freshwater Ecology* 21: 293-299.
- McDonald, D. L., T. H. Bonner, E. L. Oborny Jr., and T. M. Brandt. 2007. Effects of fluctuating temperatures and gill parasites on reproduction of the fountain darter, *Etheostoma fonticola*. *Journal of Freshwater Ecology* 22: 311-318.
- Near, T. J. and seven co-authors. 2011. Phylogeny and temporal diversification of darters (Percidae: Etheostomatinae). *Systematic Biology* 60:565-595.
- Paine, M. D., J. J. Dodson, and G. Power. 1981. Habitat and food resource partitioning among four species of darters (Percidae: Etheostoma) in a southern Ontario stream. *Canadian Journal of Zoology* 60:1635-1641.
- Phillips, C.T., M.L. Alexander, and A.M. Gonzales. 2011. Use of macrophytes for egg deposition by the endangered fountain darter. *Transactions of the American Fisheries Society* 140:1392-1397.

- Riis, T. and I. Hawes. 2002. Relationships between water level fluctuations and vegetation diversity in shallow water of New Zealand lakes. *Aquatic Botany* 74:133-148.
- Schenck J.R. and B. G. Whiteside. 1976. Distribution, habitat preference and population size estimate of *Etheostoma fonticola*. *Copeia* 4: 697-703.
- Schenck J.R., B. G. Whiteside. 1977. Reproduction, Fecundity, Sexual Dimorphism and Sex Ratio of *Etheostoma fonticola*. *American Midland naturalist*. 98: 365-375
- Schlosser, I. J. 1982. Trophic structure, reproductive success, and growth rate of fishes in a natural and modified headwater stream. *Canadian Journal of Fisheries and Aquatic Sciences* 39:968-978.
- Simon, T. P. 1999. Assessment of Balon's reproductive guilds with application to Midwestern North American freshwater fishes. 97–121. *In*: Simon, T. P. Assessing the sustainability and biological integrity of water resources using fish communities. CRC Press. New York, New York.
- Strawn, K. 1956. A method of breeding and raising three Texas darters, Part II. *Aquarium J.* 27:11-14, 17, 31.
- Tolley-Jordan, L.R. and J.M. Owen. 2008. Habitat influences snail community structure and trematode infection levels in a spring-fed river, Texas, USA. *Hydrobiologia* 600: 29-40